

Essentials

High Quality, soluble Collagen Type II (minimum 60% Collagen and 10% HA) from Chicken Cartilage.



PRODUCT PROFILE SHEET

Collagen is one of the proteins found in most connective tissues, including cartilage, bone and skin. As our body ages, it loses collagen. Other factors like diet, smoking, alcohol and sun exposure also affect collagen loss. Therefore, food supplement products can play an extremely important role.

Collagen Type II's main benefits are mainly related to joints, cartilage, and ligaments. The main, and most commonly used, source of Type II Collagen is chicken collagen, as chicken cartilage is the richest and most densely packed source of Type II Collagen.

Product Advantages

- ✓ High Quality Fine Powder
- ✓ Standardised to minimum 60% Collagen Type II
- ✓ Minimum 10% HA (Hyaluronic Acid)
- ✓ Minimum 10% Chondroitin Sulphate
- ✓ Good Solubility and Flowability
- ✓ Solvent free Extraction
- ✓ From selected approved partners
- ✓ Qualified and approved by our Regulatory team
- ✓ European Origin

Research

Orally administered hydrolysed collagen has been found to be absorbed intestinally and stimulates chondrocytes to produce type II collagen. Research suggest that hydrolysed collagen type II have the potential to repair or regenerate deteriorating collagen.^{1,2}

Approximately 50% of all the protein in the body's joint cartilage is type II collagen. The three protein components for cartilage are proteoglycan, elastin, and type II collagen.

Type II collagen fibres make up the network of bonds between elastin and proteoglycans, creating the base structure for a strong tissue. That is why several studies confirm that collagen type II supplementation may play an important role in **supporting cartilage strength**.

Collagen Type II also supports joints in absorbing the impact of movement through the day by fortifying cartilage structure and making sure that bones can move fluidly.

Collagen type II may be effective and safe in treating **knee osteoarthritis**³ and **rheumatoid arthritis**⁴, most notably in reducing pain and inflammation.

It may also **reduce inflammation and oxidative stress** (collagen II has antioxidant properties) in the human body.

Synergy Between Collagen and OptiMSM®

Collagen and Sulphur create the perfect synergy as both nutrients support the body's defences to help preserve healthy collagen in the skin.

Collagen helps in an immune-mediated remodelling of the skin⁵ while sulphur aids in the **antioxidant defence** by way of supporting **glutathione structure and function**⁶.

A study⁷ on women (aged 40-65) found that combining **OptiMSM®** with collagen improved **skin thickness, hydration, and roughness**. While collagen alone reduced roughness, the effect was significantly greater when combined with **OptiMSM®**.

The study concluded that MSM enhances collagen's benefits for skin health.

Product Range

LEHVOSS Code	Active Content	Grade	Particle Size
26002394	Min 60% Collagen Type II Min 10% Sodium Hyaluronate Min 10% Chondroitin Sulphate	Fine Powder	50 to 100 µm

Product Pack Size

Our collagen Type II (Chicken) powder is available in 25kg bags.

Product Dosage

No RDA is set for this ingredient however, up to 10 g a day is the typical use for Collagen Type II. Formulations on the market show different dosages, depending if used in combination with other ingredients.

About



LEHVOSS Nutrition is the European division of the Hamburg-based group, specialising in the sourcing, technical approval, sales and marketing of specialty ingredients for the food supplement, pharma, animal nutrition and functional foods industries.

References

1. *Desser S., Seifert J. Stimulation of type II collagen biosynthesis and secretion in bovine chondrocytes cultured with degraded collagen. Cell Tissue Res. 2003;311:393-399. doi: 10.1007/s00441-003-0702-8.*
2. *Bello A.E., Desser S. Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: A review of the literature. Med. Res. pin. 2006;22:2221-2232. doi: 10.1185/030079906X148373.*
3. *J P Lugo, Z M Saiyed, N E Lane: Efficacy and tolerability of an undenatured type II collagen supplement in modulating knee osteoarthritis symptoms, Nutrition Journal volume 15, Article number: 14 (2015), published January 2016.*
4. *W Wei, Z Ling-Ling et al. A multicenter, double-blind, randomized, controlled phase III clinical trial of chicken type II collagen in rheumatoid arthritis. Arthritis Res Ther. 2009;11(6):R180. doi: 10.1186/ar2870. Epub 2009, Dec 1.*
5. *Kalman DS & Hewlings SJ. EC Nutrition 12.11 (2018): 6840695.*
6. *Pogačnik T et al. (2023) The effect of a 12-week dietary intake of food supplements containing collagen and MSM on dermis density and other skin parameters: A double-blind, placebo-controlled, randomised four-way study comparing the efficacy of three test products. doi.org/10.1016/j.jff.2023.1058.*
7. *Barati et al. (2020) Collagen supplementation for skin health: A mechanistic systematic review. doi:10.1111/jocd.13435.*

© 2026 Any information or recommendations made for use of Seller's materials do not affect in any way Buyer's obligation to examine and/or test the Seller's goods with regard to their suitability for Buyer's purposes especially with regard to consumer use. No information given by the Seller is to be construed in any way as a guarantee regarding characteristics or duration of use, unless such information has been explicitly given as a guarantee. Any information given on the website is only applicable to the ingredients supplied by Seller and it is Buyer's obligation to ascertain how to advertise and label products containing the ingredients towards the final consumer.