

[< Back](#)

Advertise

Measurements—Again
Agilent Seahorse XF Pro Analyzer

The FASEB Journal / Volume 27, Issue S1 / p. 1076.7-1076.7

Nutrition | [Free Access](#)

A Randomized Double Blind Placebo Controlled Evaluation of MSM for Exercise Induced Discomfort/Pain.

Douglas Kalman, Samantha Feldman, Adam Samson, Diane Krieger

First published: 01 April 2013

https://doi.org/10.1096/fasebj.27.1_supplement.1076.7

Citations: 3

Abstract

Methylsulfonylmethane (MSM) has been reported to provide anti-inflammatory & antioxidant effects in mammals. Resistance exercise is known to induce both inflammation & oxidative stress resulting in muscular discomfort & pain. In a pilot-proof of concept study, we determined the effects of MSM on markers of exercise recovery & performance.

Methods

In random order 24 moderately exercise-trained men (25.5 ± 5.6 yrs) received MSM 3.0 gm/d or PLA, for 14 days, with a 17-day washout between. The study included 3 tests: baseline, no product & the 2nd & 3rd following 14d supplementation with MSM & PLA. Each test consisted of 2 visits. At the 1st visit, subjects performed stressing exercise; 28 total sets of leg extensions, sets 1–25, predetermined weight, 10 reps each, sets 26–28 to muscular failure at 70% 1-RM (performance). At the 2nd visit (48 hrs later), subjects performed 12 total sets of leg ext, sets 1–9, predetermined weight, sets 10–12 to muscular failure, 70% 1-RM (performance). Muscle discomfort/pain (10-point VAS scale), inflammation (hs-CRP & IL-6), blood antioxidant status (TEAC & SOD), & homocysteine were measured before the stressing exercise and 2 & 48 hours post exercise. Exercise performance was also measured following 14d supplementation with MSM/PLA (sets 26–28 and sets 10–12). In order to eliminate sequence effects, results are provided on the 1st product as compared to baseline (comparison of the 1st & 2nd tests).

[Back](#)

MSM intervention resulted in significantly less pain/discomfort vs. PLA from baseline to 2 hrs (1.55±0.82 vs 3.75±2.58, p=0.012). Change in IL-6 was significant within the MSM & PLA & between MSM (0.54±0.76) vs PLA (−0.58±0.97) p = 0.006. There were no significant differences between MSM & PLA noted for the other biomarkers.

Conclusion

MSM may help alleviate the discomfort/pain that can follow a stressful exercise session.

Citing Literature



© 2024 Federation of American Societies for Experimental Biology (FASEB)

ABOUT WILEY ONLINE LIBRARY

- Privacy Policy
- Terms of Use
- About Cookies
- Manage Cookies
- Accessibility
- Wiley Research DE&I Statement and Publishing Policies

HELP & SUPPORT

- Contact Us
- Training and Support
- DMCA & Reporting Piracy

OPPORTUNITIES

- Subscription Agents
- Advertisers & Corporate Partners

CONNECT WITH WILEY

- The Wiley Network
- Wiley Press Room

< Back

and training of artificial technologies or similar technologies.

WILEY